

**Oral Surgery
Post Operative Care**



Dr. Allan J. Libunao
Dr. Richard H. Kim

580 East Boughton Road
Suite B
Bolingbrook, IL 60440
Office (630) 972-1599
Fax (630) 972-1050

24 hour answering service

Bleeding

Bite down on gauze for 1 hour, or until bleeding has stopped. Extra gauze will be provided if bleeding resumes. Oozing of blood is normal for the first 24 hours. Excessive bleeding can be controlled by placing gauze or tea bag over the wound and biting firmly for 30 minutes. Avoid frequent pack changes.

DO NOT spit, rinse, or use a straw for the next 24 hours. After 24 hours, rinse mouth with warm salt water after every meal for the first week. Brush your teeth as usual, but stay away from the areas of surgery. Pain in the area, the ear, adjacent teeth, difficulty in swallowing and opening the mouth, alterations in your bite, swelling and bruising may occur and are quite normal.

Swelling

Your swelling will take 48 hours to reach its maximum and then will slowly subside. To minimize the swelling, apply ice to the side of your face at 20 minute intervals for the first 48 hours. After the third day, if swelling is still present, use moist heat instead of ice.

Diet

Stay on a soft food diet for the first couple days (i.e. soups, juices, milk shakes, mash potatoes, etc.) Good nutrition is necessary for normal healing.

Smoking

Limit or avoid smoking as long as possible after extractions, as it tends to disturb the healing and may cause a dry socket.

Medications

Take and finish your prescription for antibiotics (i.e. penicillin, amoxicillin) as directed. Antibiotic medication may decrease the effectiveness of birth control pills. Take your first dosage of the prescribed pain medication before the local anesthesia wears off. The local will last about 3 hours. Do not take the prescription pain medication on an empty stomach. Do not drive or operate heavy machinery while take the prescription pain medication. Avoid alcoholic beverages while taking prescription pain medication. For minor discomfort, you can take two tablets of acetaminophen (Tylenol) or ibuprofen (Advil) every 4 hours.

**IN CASE OF ANY UNUSUAL OCCURRENCES,
PLEASE CALL OUR OFFICE 24-HOUR EMERGENCY
SERVICE
(630) 972-1599**